

Study of Risk Factors in Stroke in Tertiary Level Hospital of Pokhara: A Cross-Sectional Study

Roshan Yadav¹, Ajit Kumar Sah², Sneha Shah³, Archana Shah⁴, Sabita Shah⁵, Priti Yadav⁶

¹Gandaki Medical College and Teaching Hospital, Pokhara

²Patan Academy of Health Sciences- School of Medicine, Kathmandu, Nepal

³Nobel Medical College Teaching Hospital, Biratnagar, Nepal

⁴Central Department of Public Health, Tribhuvan University, Kathmandu, Nepal

⁵Kist Medical College Teaching Hospital, Kathmandu, Nepal

⁶Kist Medical College Teaching Hospital, Kathmandu, Nepal

Email address: doc.ajitks@gmail.com

Abstract— Background: Stroke is defined as an "acute neurologic dysfunction of vascular origin with sudden (within seconds) or at least rapid (within hours) occurrence of symptoms and signs corresponding to the involvement of focal areas in the brain. The study aims to study the various risk factors in stroke. **Materials and Methods:** This was a hospital-based cross-sectional study conducted on 200 patients having the problems of stroke in Gandaki Medical College and Teaching Hospital, Pokhara from August 2020 to September 2021. Data of all the patients were collected and analyzed using the appropriate statistical tool. **Results:** A total of 200 patients with stroke were recruited; patients each had an ischemic and hemorrhagic stroke. Demographic, as well as clinical characteristics were recorded and data were analyzed by using statistical software like SPSS. **Conclusion:** There was a statistically significant association of risk factors of stroke and age in our study having a p -value <0.005 . There was a statistically significant association between hypertension and stroke $p=0.002$.

Keywords— Stroke, risk, hemorrhagic, hypertension, thrombosis.

I. INTRODUCTION

Stroke is defined as an "acute neurologic dysfunction of vascular origin with sudden (within seconds) or at least rapid (within hours) occurrence of symptoms and signs corresponding to the involvement of focal areas in the brain"¹. It is a sudden onset of a very non-convulsive focal neurological defect that can last for 1 day. Stroke is the 3rd leading cause accounting for 10% of all death in the world². The cerebrovascular accident is a worldwide health problem. It is the third most common cause of death in the developing world and hence is an important cause of morbidity, mortality, and disability in developed as well as developing countries. Although there are substantial differences in frequencies from place to place, cerebral thrombosis is usually the most frequent form of stroke encountered in clinical studies followed by cerebral hemorrhage, subarachnoid hemorrhage, and cerebral embolism. Stroke management differs from patient to patient depending on the type of stroke, associated risk factors, and presence of complications³. The management of hemorrhagic stroke differs from that of ischemic stroke, though the etiological factors are the same. The advent of investigative facilities like CT scan, MRI, digital subtraction angiography, etc., has brought into focus the discrepancies between clinical

and radiological diagnosis. Though a good clinician is most often accurate in differentiating ischemic from hemorrhagic stroke, sometimes misdiagnosis may result in therapeutic disasters in this era of consumer activism, there is a crying need to prevent such errors of diagnosis⁴. It is a major global health issue that prematurely claims millions of otherwise healthy and productive lives each year. The WHO predicts that the total stroke burden will increase from approximately 38 million Disability Adjusted Life Years (DALYs) in 1991 to 51 million (DALYs) in 2020⁵.

Strokes are rare in younger people and become increasingly common with older age but at the age of 45 years, the chances of having a stroke in the next 20 years are 1 in 30⁶. There has been a definite increase in the prevalence and incidence of stroke disorder in India over the last 30 years⁷.

Stroke represented 1.2% of total deaths in India⁸. The average annual incidence rate of stroke in India currently is 145 per 100,000 populations, which is higher than the western nations and Indians may also be genetically prone for stroke due to the high prevalence of metabolic syndrome⁹. Ischemic and hemorrhagic stroke accounts for approximately 85% and 15%, respectively¹⁰. Stroke is a clinical culmination of many complex processes and interacting pathways, none of which is individually necessary or sufficient to cause the outcome. There is considerable evidence from twin studies and numerous epidemiologic studies have documented that stroke has a significant genetic component¹¹. Studies looking at sibling risk and/or offspring showed significant risk in first-degree relatives of stroke patients, with odds ratios that average 1.76¹². Family studies have shown an association of hemorrhagic stroke and a positive family history of hemorrhagic stroke, indicating an underlying genetic etiology¹³.

The cause of most common strokes is multifactorial and involves both genetic variants, environmental triggers such as smoking or diet, and most likely results from that of additive or multiplicative effect of a wide spectrum of pathogenic alleles (a gene dose effect), each of which conferring a small degree of risk but the identification of individual causative mutations remains problematic¹⁴. Many common risk factors for stroke like diabetes and arterial hypertension are partly

inherited; so many genetic loci contribute more or less to the stroke phenotype¹⁵.

Risk factors for hemorrhagic stroke have become evident, including heavy alcohol intake and anticoagulant treatment. Less frequent and convincingly proven risk factors are cocaine use and low serum cholesterol¹⁶.

Many candidate genes for hypertension have been suggested including genes for rennin angiotensin-converting enzyme, aldosterone, calmodulin, ion ATPases, phospholipase, kallikrein, endothelin, and androgenic receptors. Genome-wide linkage surveys have found linkage for blood-pressure loci on almost all chromosomes¹⁷.

Several genetic polymorphisms have been also associated with an increased risk for apparently sporadic hemorrhagic stroke. The influence of *APOE* genotype on sporadic cerebral amyloid angiopathy-related –hemorrhagic stroke has been studied intensively, with contradicting results like in ischemic stroke. It is now accepted that the *APOE* E2 allele predisposes to rupture of β A4 laden blood vessels, thus increasing the chance of sporadic cerebral amyloid angiopathy - related-hemorrhagic stroke¹⁸.

Atherosclerosis is a well-known risk factor for stroke and the genetic variants of the low-density lipoprotein receptor have already been highly informative for the genetics of atherosclerosis in vascular disease¹⁹. Ischemic stroke can be caused by many monogenic disorders and in such cases, stroke is frequently part of a multisystem disorder²⁰. The ischemic stroke itself consists of many different phenotypes which may each have different genetic profiles²¹.

A genome-wide linkage study on Icelandic individuals with stroke has indicated that a gene on chromosome 5 may contribute significantly to the risk of stroke²². In many of the studies, the risk persisted with hypertension, hyperlipidemia, diabetes, and smoking. These studies support the notion that stroke, as with all common diseases, occurs at the interface between genes and the environment. However, the significant genetic component to stroke risk is not fully explained by the genetic components driving the known medical risk factors²³.

To relieve the burden of stroke, we need to understand the mechanisms that will form the basis of improved prevention and treatment. In the post-genomic era, multiple genes have been termed low-penetrance genes and various studies have focused on different variants of these genes known as polymorphisms. Three strategies are being used to find stroke genes: positional cloning applied to rare monogenic forms of stroke, candidate gene association studies of common stroke, and positional cloning applied to common stroke²⁴.

There may be inter-individual variation in the studied subjects. Extensive effort has been expended to identify the association between single nucleotide polymorphisms (SNPs) in functional candidate genes and stroke²⁵. However, the candidate gene approach is limited by how much is the underlying mechanism of the disease being investigated. In the present study, we have employed targeted candidate gene analysis to look for the association of genetic variants with the risk of stroke and their clinical subtypes. Candidate genes were chosen a priori based on assumed biological relevance.

Cerebrovascular accident is a worldwide health problem. It is the third most common cause of death in the developing world and hence is an important cause of morbidity, mortality, and disability in developed as well as developing countries. Although there are substantial differences in frequency from place to place, cerebral thrombosis is usually the most frequent form of stroke encountered in clinical studies followed by cerebral hemorrhage, subarachnoid hemorrhage, and Cerebral Embolism.

Stroke management differs from patient to patient depending on the type of stroke, associated risk factors, and presence of complications. The management of Hemorrhagic stroke differs from that of Ischaemic stroke, though the aetiological factors are the same. The advent of investigative facilities like CT scan, MRI, Digital subtraction angiography, etc., has brought into focus the discrepancies between clinical and radiological diagnosis. Though a good clinician is most often accurate in Differentiating ischaemic from hemorrhagic stroke, sometimes misdiagnosis may result in therapeutic disasters. In this era of Consumer activism, there is a crying need to prevent such errors of diagnosis²⁶⁻³⁰. Very limited information is available on the genetic basis of stroke especially in hemorrhagic stroke of Nepali population. In the present study we explored the possible risk factors of Nepali population.

II. METHODS

Subjects

The study population consisted of 200 both ischemic and hemorrhagic stroke patients. Based on inclusion/exclusion criteria, patients were recruited from the Neurology outpatient and inpatient of Gandaki Medical College Hospital. Also, question regarding the language they speak, the food they eat, determined the ethnicity. The study was ethically approved by institutional ethics committee.

Research Design

Hospital based cross-section observational study.

Type of Design

Hospital based cross-section observational study.

Research Question

To observe the risk factor according to age

To see the risk factor according to sex

Inclusion criteria:

1. Age above 35 years
2. Brain CT-scan indicative of cerebral infarction or intra-cerebral hemorrhage
3. MRI indicative of cerebral infarction or intra-cerebral hemorrhage

Exclusion criteria: The patients with following disorders were excluded

1. Patients in whom the cerebral hemorrhage secondary to trauma or previous coagulation disorders

Study Duration

The study was conducted from July 2020 to August 2021

Study Population

Study population was any patients fulfilling the inclusion criteria with clinical features of stroke in Gandaki Medical College, Pokhara, Nepal.

Sampling Method

The sampling technique that was employed was non-probability convenience technique.

Sampling Technique: The participants that are required for the research are 200. As per the study of literature of prevalence of hypertension in stroke patients was 47.2%

$$n = Z^2 p^* (1-p) / d^2$$

$$n = (1.962 \times 47.2 \times 52.8 / 7.08)^2$$

$$n = 191.01$$

$$= 192 \text{ where, } n = \text{the desired samples men}$$

$$z = \text{the standard normal deviate usually set at 1.96 (or more)}$$

$$p = \text{Prevalence at 2.0 in previous study}$$

$$Q-1.0-P$$

$$d = \text{Absolute error or precision (15\%)}$$

$$\text{i.e. 15\% of 47.2 which is 7.08}$$

For convenience 200 samples were taken into consideration.

Tools and technique

Data management and statistical analysis statistical methods

The data obtained by history, clinical examination and on investigations as per case entered in Microsoft excel. Demographic as well as clinical characteristics were recorded and data were analyzed by using statistical software like SPSS 15.

III. RESULTS

In the present case-control study, we selected 200 patients having hemorrhage and the same number had ischemic stroke. The risk factors of stroke such as age, gender, smoking, tobacco, alcohol intake, hypertension (systolic blood pressure >140 mm Hg or diastolic blood pressure >90 mm Hg, history of hypertension or on antihypertensive drugs), diabetes (fasting blood sugar > 106 mg/dl, 2 hr postprandial blood sugar > 200 mg/dl, or on oral hypoglycemics/insulin), dyslipidemia (defined as a history of an abnormal lipid profile that required either dietary or pharmacological intervention, or an abnormal lipid profile) were recorded. History of heart disease such as atrial fibrillation, rheumatic heart disease or coronary artery disease, family history of hypertension, diabetes mellitus and stroke in first degree relatives were enquired.

3.1 Distribution of patients according to sex

The majority of the stroke patients were male i.e. 120 (60%) in the study group while the rest were females i.e. 40 %.

TABLE 1: Distribution of patients according to sex

N=200

	(n)	Percentage (%)
Male	120	60
Females	80	40

3.2 Distribution of patients according to age group

The persons are found to be highest in the age group of 30-39 i.e., 40, whereas the least number is found in the age group of 90-99.

TABLE 2: Distribution of cases according to age

N=200

Age in years	No. of patients	Percentage (%)
20-29	33	16.5
30-39	40	20.0
40-49	35	17.5
50-59	32	16.0
60-69	25	12.5
70-79	20	10.0
80-89	14	7.0
90-99	1	0.5

3.3 Age and gender of study groups

Mean age of stroke patients with healthy controls were compared using an independent t-test. The mean age of healthy controls was 55.25 years (SD±10.65) while ischemic stroke patients had mean age of 53.84 years (SD±12.14) and hemorrhagic stroke had 56.82 years (SD±10.35). The two-sided P value obtained on comparing mean ages of stroke patients with healthy controls was not significant (Ischemic stroke, P=0.32; hemorrhagic stroke, P=0.22). In healthy control group, there were 120 male and 80 female.

3.4 Demographic and risk factors:

109 persons visiting the hospital had clinical features related to stroke. Most patients were suffering from hemiparesis i.e. 84(77.06%) whereas 3(2.75%) persons were suffering from headache and vomiting.

TABLE 3: Distribution of clinical features

N=109

Clinical Features	(n)	Percentage (%)
Hemiparesis / hemiplegia	84	77.06
7 th nerve palsy	42	38.5
Speech disorder	35	32.11
Comatose	19	17.43
Convulsions	4	3.66
Headache & Vomiting	3	2.75

3.5 Distribution of Stroke subtypes

Among 109 patients, most of them i.e. 69.72% had the cerebral infarction as subtype of stroke while only 2.75% were categorized under subarachnoid hemorrhage subtype.

TABLE 4: Distribution of Stroke subtypes

N=109

Stroke Subtypes	(n)	Percentage
Cerebral infarction	76	69.72
Intracerebral hemorrhage	16	14.67
Subarachnoid hemorrhage	3	2.75
Undetermined	14	12.84

3.6 Demographic and clinical characteristics of study subjects

Among the patient group, most of them i.e. 37.5% were hypertensive, 7.5% were diabetic and Ischemic Heart Disease being the least i.e. 0.5%. Hypertension was found to be the main cause of stroke. In the patient group, 5% were smokers, 9.4% chewed tobacco and 50% alcohol consumption patients showed more risk of stroke.

3.7 Stroke outcome at 30 days

Out of the total, nearly half of them i.e. 47.71% had stable conditions at 30 days while 28.85% had improved outcomes.

TABLE 5: Distribution of Risk Factors in Stroke

N=200

Risk Factors	(n)	Percentage (%)
Hypertension	75	37.5
Dyslipidemia	32	16.0
H/O Stroke or Transient Ischemic Attack (TIA)	24	12.0
H/O hypertension	28	14.0
Diabetes mellitus	15	7.5
Valvular heart disease	25	12.5
Ischemic Heart Disease	1	0.5

TABLE 6: Stroke Outcome at 30 days

N=109

Outcomes	(n)	Percentage (%)
Stable	52	47.71%
Deteriorated	31	28.44%
Improved	26	23.85%

IV. DISCUSSION

Stroke is a common neurological disease and a leading cause of severe disability and death. Stroke is believed to be a multi-factorial disorder or complex trait for which it is not possible to demonstrate classical patterns of inheritance; however, twin, family and animal model studies have consistently suggested a genetic influence on stroke risk and prognosis⁸⁹.

Diagnosed hemorrhagic cases on CT scan and MR angiography were done among the patients who were normal tensile or had lobar hemorrhages. Therefore, other causes of hemorrhagic stroke such as arteriovenous malformation, aneurysm, and tumor bleeding were unlikely. Deep brain hemorrhage is mainly due to hypertension and is more common in Asian countries whereas lobar hemorrhagic stroke in the elderly is due to cerebral amyloid angiopathy and may have a different pathophysiological basis.

However, according to some studies, the frequency and contribution of hypertension to Intra-Cerebral Hemorrhage (ICH) were similar in patients with lobar hemorrhage compared with deep hemispheric, cerebellar, or pontine hemorrhage⁹⁰⁻⁹³. Patients with documented Cerebral Amyloid Angiopathy (CAA) may also have hypertensive angiopathy in 30% of patients. Hypertension may increase the tendency of hypertension-related hemorrhage and vice versa⁹⁴.

Hypertension was found to be the most important risk factor for ICH. In this study, Hypertension was detected in 80% of cases which correlated well with other studies. Probably other risk factors like smoking 60% and alcoholism (50%) were also responsible to the occurrence of ICH in males. In addition to hypertension, the male community is more prone to indulging in smoking and alcohol consumption when compared to females in developing countries like Nepal. Excessive intake of alcohol might result in increased Very Low Density Lipoprotein (VLDL) and Low Density Lipoprotein (LDL) content resulting in accelerated atherosclerosis. Diabetes mellitus was noted in 12% of cases in Archana Verma et al., 95.14% in Sk Mohapatra et al.,⁹⁶ and 8.57% in Akbari et al.⁹⁷, and 22% in the present study. Clinical features like headache in 68%, vomiting in 58%, and convulsions in 16% of cases were important presenting features, which enabled a clinical diagnosis of ICH in this

study. The onset of stroke was also taken into consideration. The diagnosis of ICH was considered with sudden onset of illness. In 200 cases, the onset is sudden i.e. 74%. Hemiplegia (80 cases) 80% was the most common presentation in the present study. Headache was noted as the most common presenting symptom and Hemiplegia was the most initial neurological sign in this study.

In the present study, a clinical diagnosis of ICH was made in 148 cases (74%) and the remaining 52 cases (26%) were considered to be Ischemic strokes. However, a CT scan showed ICH in all 200 cases. Thus, there is a discrepancy of 26% under-diagnosis. This significant number of misdiagnosis/under-diagnosis and therapeutic problems associated with misdiagnosis/under-diagnosis strongly favors and recommends the routine use of a CT Scan Brain for establishing the final diagnosis in all suspected cases of Cerebrovascular Accidents (CVAs). Size of Haematoma, site of hemorrhage, and Associated Mass Effect on CT scan were the important prognostic factors observed in this study. The site of hemorrhage also plays a vital role in predicting the outcome in patients with Intracerebral Haemorrhage. Mortality was lower in patients with capsule-ganglionic hematoma (21.4%). The level of sensorium at the time of onset is the most important prognostic factor of the final outcome. In the present study, most patients presented with coma, fewer patients presented with altered sensorium, and two Patient was conscious at the time of admission.

Amit Kumar et.al⁹⁸ dealt with 100 patients of stroke who were admitted to B. R. D. Medical College, Gorakhpur, India. Each patient was analyzed in detail about clinical presentation and the investigations were aimed to establish the pathologic type of stroke and estimation of risk factors. Stroke incidence was more in males (Male: Female= 1.43:1). Maximum incidence of stroke was in the 6th decade (32%) followed by the 7th decade (30%). Among modifiable risk factors, a history of hypertension was the commonest (51%) followed by smoking (36% of patients) exclusively, found in males. Hemiparesis was the most common presentation (95%) followed by altered sensorium (55%). Chest X-ray was abnormal in 16% of patients, abnormal ECG was found in 27% of patients and abnormal lipid values were found in 54 patients. They concluded that apart from control of hypertension and diabetes, abnormal lipid profile remains an important modifiable risk factor for stroke.

V. CONCLUSION

The majority of the stroke patients were male i.e. 120 (60%) in the study group and hypertension was found to be the main cause of stroke. 55.40% of the ischemic stroke patients were under the large vessel disease subgroup. In the patient group 5% were smokers, 9.4% chewed tobacco and 50% alcohol consumption patients showed more risk of stroke. Family H/o stroke was present in 15.54% of the total patients. History of TIA/stroke was observed in 11.65% of the total study subjects showed a higher (twentyfold) risk in lobar.

REFERENCES

1. Abraham J, Rao PS, Inbaraj SG, Shetty G, Jose CJ. An epidemiological study of hemiplegia due to stroke in South India. *Stroke*. 1970 Nov-Dec;1(6):477-81.
2. Achrol AS, Pawlikowska L, McCulloch CE, Poon KY, Ha C, Zaroff JG, et al. Tumor necrosis factor-alpha-238G>A promoter polymorphism is associated with increased risk of new hemorrhage in the natural course of patients with brain arteriovenous malformations. *Stroke*. 2006 Jan;37(1):231-4.
3. Adams HP, Jr., Bendixen BH, Kappelle LJ, Biller J, Love BB, Gordon DL, et al. Classification of subtype of acute ischemic stroke. Definitions for use in a multicenter clinical trial. TOAST. Trial of Org 10172 in Acute Stroke Treatment. *Stroke*. 1993 Jan;24(1):35-41.
4. Aikawa M, Rabkin E, Okada Y, Voglic SJ, Clinton SK, Brinckerhoff CE, et al. Lipid lowering by diet reduces matrix metalloproteinase activity and increases collagen content of rabbit atheroma: a potential mechanism of lesion stabilization. *Circulation*. 1998 Jun 23;97(24):2433-44.
5. Al-Allawi NA, Avo AS, Jubrael JM. Methylenetetrahydrofolate reductase C677T polymorphism in Iraqi patients with ischemic stroke. *Neurol India*. 2009 Sep-Oct;57(5):631-5.
6. Alluri RV, Mohan V, Komandur S, Chawda K, Chaudhuri JR, Hasan Q. MTHFR C677T gene mutation as a risk factor for arterial stroke: a hospital based study. *Eur J Neurol*. 2005 Jan;12(1):40-4.
7. Anderson CD, Biffi A, Rost NS, Cortellini L, Furie KL, Rosand J. Chromosome 9p21 in ischemic stroke: population structure and meta-analysis. *Stroke*. 2010 Jun;41(6):1123-31.
8. Antikainen R, Jousilahti P, Tuomilehto J. Systolic blood pressure, isolated systolic hypertension and risk of coronary heart disease, strokes, cardiovascular disease and all-cause mortality in the middle-aged population. *Journal of Hypertension*. 1998;16(5):577-83.
9. Ariesen MJ, Claus SP, Rinkel GJ, Algra A. Risk factors for intracerebral hemorrhage in the general population: a systematic review. *Stroke*. 2003 Aug;34(8):2060-5.
10. Ariyaratnam R, Casas JP, Whittaker J, Smeeth L, Hingorani AD, Sharma P. Genetics of ischaemic stroke among persons of non-European descent: a meta-analysis of eight genes involving approximately 32,500 individuals. *PLoS Med*. 2007 Apr;4(4):e131.
11. Arruda VR, Siqueira LH, Goncalves MS, von Zuben PM, Soares MC, Menezes R, et al. Prevalence of the mutation C677 --> T in the methylene tetrahydrofolate reductase gene among distinct ethnic groups in Brazil. *Am J Med Genet*. 1998 Jul 24;78(4):332-5.
12. Bader MK, & Littlejohns, L. R. . (Eds) (2004) St Louis: AANN core curriculum for neuroscience nursing (4th ed) Elsevier. 2004.
13. Baillie G, MacKenzie SJ, Houslay MD. Phorbol 12-myristate 13-acetate triggers the protein kinase A-mediated phosphorylation and activation of the PDE4D5 cAMP phosphodiesterase in human aortic smooth muscle cells through a route involving extracellular signal regulated kinase (ERK). *Mol Pharmacol*. 2001 Nov;60(5):1100-11.
14. Bak S, Gaist D, Sindrup SH, Skytthe A, Christensen K. Genetic liability in stroke: a long-term follow-up study of Danish twins. *Stroke*. 2002 Mar;33(3):769-74.
15. Bamford J, Sandercock P, Dennis M, Burn J, Warlow C. A prospective study of acute cerebrovascular disease in the community: the Oxfordshire Community Stroke Project-- 1981-86.
16. Bamford J, Sandercock P, Dennis M, Burn J, Warlow C. Classification and natural history of clinically identifiable subtypes of cerebral infarction. *Lancet*. 1991 Jun 22; 337(8756):1521-6.
17. Banerjee I, Gupta V, Ganesh S. Association of gene polymorphism with genetic susceptibility to stroke in Asian populations: a meta-analysis. *J Hum Genet*. 2007; 52(3):205-19.
18. Banerjee I, Gupta V, Ahmed T, Faizaan M, Agarwal P, Ganesh S. Inflammatory system gene polymorphism and the risk of stroke: a case-control study in an Indian population. *Brain Res Bull*. 2008 Jan 31; 75(1):158-65.
18. Banerjee TK, Das SK. Epidemiology of stroke in India. *Neurology Asia*. 2006; 11:1-4.
19. Banerjee TK, Mukherjee CS, Sarkhel A. Stroke in the urban population of Calcutta--an epidemiological study. *Neuroepidemiology*. 2001 Aug;20(3):201-7.
20. Banerjee TK, Roy MK, Bhoi KK. Is stroke increasing in India--preventive measures that need to be implemented. *J Indian Med Assoc*. 2005 Mar;103(3):162, 4, 6 passim.
21. Bansal BC, Prakash C, Jain AL, Brahmanandam KR. Cerebrovascular disease in young individuals below the age of 40 years. *Neurol India*. 1973 Mar;21(1):11-8.
22. Batalla A, Alvarez R, Reguero JR, Hevia S, Iglesias-Cubero G, Alvarez V, et al. Synergistic effect between apolipoprotein E and angiotensinogen gene polymorphisms in the risk for early myocardial infarction. *Clin Chem*. 2000 Dec;46(12):1910-5.
23. Bevan S, Porteous L, Sitzer M, Markus HS. Phosphodiesterase 4D gene, ischemic stroke, and asymptomatic carotid atherosclerosis. *Stroke*. 2005 May;36(5):949-53.
24. Bevan S, Dichgans M, Wiechmann HE, Gschwendtner A, Meitinger T, Markus HS. Genetic variation in members of the leukotriene biosynthesis pathway confer an increased risk of ischemic stroke: a replication study in two independent populations. *Stroke*. 2008 Apr;39(4):1109-14.
25. Bhattacharya S, Saha SP, Basu A, Das SK. A 5 years prospective study of incidence, morbidity and mortality profile of stroke in a rural community of eastern India. *J Indian Med Assoc*. 2005 Dec;103(12):655-9.
26. Bianchi G, Ferrari P, Staessen JA. Adducin polymorphism: detection and impact on hypertension and related disorders. *Hypertension*. 2005 Mar;45(3):331-40.
27. Birk S, Edvinsson L, Olesen J, Kruuse C. Analysis of the effects of phosphodiesterase type 3 and 4 inhibitors in cerebral arteries. *Eur J Pharmacol*. 2004 Apr 5;489(1-2):93-100.
28. Biswas A, Ranjan R, Meena A, Akhter MS, Yadav BK, Munisamy M, et al. Homocysteine levels, polymorphisms and the risk of ischemic stroke in young Asian Indians. *J Stroke Cerebrovasc Dis*. 2009 Mar-Apr;18(2):103-10.
29. Blank-Reid C. How to have a stroke at an early age: the effects of crack, cocaine and other illicit drugs. *J Neurosci Nurs*. 1996 Feb;28(1):19-27.
30. Bonita R, Broad JB, Beaglehole R. Changes in stroke incidence and case-fatality in Auckland, New Zealand, 1981-91. *Lancet*. 1993 Dec 11;342(8885):1470-3.
31. Bonita R, Solomon N, Broad JB. Prevalence of stroke and stroke-related disability. Estimates from the Auckland stroke studies. *Stroke*. 1997 Oct;28(10):1898-902.
32. Brass LM, Alberts MJ. The genetics of cerebrovascular disease. *Baillieres Clin Neurol*. 1995 Aug;4(2):221-45.
33. Brass LM, Isaacsohn JL, Merikangas KR, Robinette CD. A study of twins and stroke. *Stroke*. 1992 Feb;23(2):221-3.
34. Brattstrom L, Wilcken DE, Ohrvik J, Brudin L. Common methylenetetrahydrofolate reductase gene mutation leads to hyperhomocysteinemia but not to vascular disease: the result of a meta-analysis. *Circulation*. 1998 Dec 8;98(23):2520-6.
35. Broderick J, Brott T, Tomsick T, Leach A. Lobar hemorrhage in the elderly. The undiminishing importance of hypertension. *Stroke*. 1993 Jan;24(1):49-51.
36. Brophy VH, Ro SK, Rhee BK, Lui LY, Lee JM, Umblas N, et al. Association of phosphodiesterase 4D polymorphisms with ischemic stroke in a US population stratified by hypertension status. *Stroke*. 2006 Jun;37(6):1385-90.
37. Brown RD, Whisnant JP, Sicks JD, O'Fallon WM, Wiebers DO. Stroke incidence, prevalence, and survival: secular trends in Rochester, Minnesota, through 1989. *Stroke*. 1996 Mar;27(3):373-80.
38. Bruno A, Biller J, Adams HP, Jr., Clarke WR, Woolson RF, Williams LS, et al. Acute blood glucose level and outcome from ischemic stroke. Trial of ORG 10172 in Acute Stroke Treatment (TOAST) Investigators. *Neurology*. 1999 Jan 15;52(2):280-4.
39. Bullock R, Mendelow AD, Teasdale GM, Graham DI. Intracranial haemorrhage induced at arterial pressure in the rat. Part 1: Description of technique, ICP changes and neuropathological findings. *Neurol Res*. 1984 Dec;6(4):184-8.
40. Burchfiel CM, Curb JD, Rodriguez BL, Abbott RD, Chiu D, Yano K. Glucose intolerance and 22-year stroke incidence. The Honolulu Heart Program. *Stroke*. 1994 May;25(5):951-7.
41. Byth BC, Billingsley GD, Cox DW. Physical and genetic mapping of the serpin gene cluster at 14q32.1: allelic association and a unique haplotype associated with alpha 1-antitrypsin deficiency. *Am J Hum Genet*. 1994 Jul;55(1):126-33.
42. Cambien F, Poirier O, Lecerf L, Evans A, Cambou JP, Arveiler D, et al. Deletion polymorphism in the gene for angiotensin-converting enzyme

- is a potent risk factor for myocardial infarction. *Nature*. 1992 Oct 15;359(6396):641-4.
43. Casas JP, Hingorani AD, Bautista LE, Sharma P. Meta-analysis of genetic studies in ischemic stroke: thirty-two genes involving approximately 18,000 cases and 58,000 controls. *Arch Neurol*. 2004 Nov;61(11):1652-61.
 44. Castellano M, Muiesan ML, Rizzoni D, Beschi M, Pasini G, Cinelli A, et al. Angiotensin-converting enzyme I/D polymorphism and arterial wall thickness in a general population. The Vobarno Study. *Circulation*. 1995 Jun 1;91(11):2721-4.
 45. Chamorro A, Revilla M, Obach V, Vargas M, Planas AM. The -174G/C polymorphism of the interleukin 6 gene is a hallmark of lacunar stroke and not other ischemic stroke phenotypes. *Cerebrovasc Dis*. 2005;19(2):91-5.
 46. Chen CH, Lo YK, Ke D, Liu CK, Liou CW, Wu HL, et al. Platelet glycoprotein Ia C807T, Ib C3550T, and IIIa Pl(A1/A2) polymorphisms and ischemic stroke in young Taiwanese. *J Neurol Sci*. 2004 Dec 15;227(1):1-5.
 47. Chen CH, Eng HL, Chang CJ, Tsai TT, Lai ML, Chen HY, et al. 4G/5G promoter polymorphism of plasminogen activator inhibitor-1, lipid profiles, and ischemic stroke. *J Lab Clin Med*. 2003 Aug;142(2):100-5.
 48. Chen YC, Lee-Chen GJ, Wu YR, Hu FJ, Wu HC, Kuo HC, et al. Analyses of interaction effect between apolipoprotein E polymorphism and alcohol use as well as cholesterol concentrations on spontaneous deep intracerebral hemorrhage in the Taiwan population. *Clin Chim Acta*. 2009 Oct;408(1-2):128-32.
 49. Choi BO, Kim NK, Kim SH, Kang MS, Lee S, Ahn JY, et al. Homozygous C677T mutation in the MTHFR gene as an independent risk factor for multiple small-artery occlusions. *Thromb Res*. 2003;111(1-2):39-44.
 50. Chowdhury AH, Yokoyama T, Kokubo Y, Zaman MM, Haque A, Tanaka H. Apolipoprotein E genetic polymorphism and stroke subtypes in a Bangladeshi hospital-based study. *J Epidemiol*. 2001 May;11(3):131-8.
 51. Collins R, MacMahon S. Blood pressure, antihypertensive drug treatment and the risks of stroke and of coronary heart disease. *Br Med Bull*. 1994 Apr;50(2):272-98.
 52. Cronin S, Furie KL, Kelly PJ. Dose-related association of MTHFR 677T allele with risk of ischemic stroke: evidence from a cumulative meta-analysis. *Stroke*. 2005 Jul;36(7):1581-7.
 53. Kasper, Braunwald, Fauci.; *Harrisons principles of Internal Medicine* 18th edn, Vol -2 :364;2513-35.
 54. Hauser, Longo, Jameson; *Harrisons principles of Internal Medicine* 17th edn, Vol -2 :364;2513-35.
 55. Heyman A et al : *The Pilot Stroke Data Bank: definition, design and data*. *Stroke* 1984; 15:740.
 56. Tomsick T et al: The risk of subarachnoid and intracerebral haemorrhages in blacks as compared with whites. *N Engl J Med* 1992;326:733.
 57. Cheyne J. *Cases of apoplexy and lethargy with observations on comatose patients*. London:Thomas Underwood, 1812.
 58. Gowers W.A *Manual of Diseases of the Nervous System*, Vol - 2.(2nd ed). London: J&A Churchill, 1892;3:84-421.
 59. Osler W. *The Principles and Practices of Medicine* (5th ed). New York: Appleton, 1903;997- 1008.
 60. Airing C, Merritt H. Differential diagnosis between cerebral hemorrhage and cerebral thrombosis: clinical and pathological study of 245 cases. *Arch Internal Med* 1935;56:435-456.
 61. Kase CS, Caplan LR, *Intracerebral haemorrhage* Boston: Butterworth - Heinemann, 1994.
 62. Kunitz SC, Gross CR : *The Pilot Stroke Data Bank: definition, design and data*. *Stroke*, 1984; 15:740.
 63. Broderick J, Brott T : The risk of subarachnoid and intracerebral haemorrhages in blacks as compared with whites. *N Engl J Med*, 1992; 326:733.
 64. Sacco RL, Wolf PA et al: Subarachnoid and intracerebral haemorrhage: natural history, prognosis and precursive factors in the Framingham Study. *Neurology* (NY), 1984; 34:847.
 65. Bharucha NE et al: Subarachnoid and intracerebral haemorrhage: natural history, prognosis and precursive factors in the Framingham Study. *Neurology* (NY), 1984; 34:847.
 66. I-Iertzberg V: Hypertension as a risk factor for spontaneous intracerebral haemorrhage. *Stroke*, 1986; 17:1078.
 67. Whisnant JP: Primary intracerebral haemorrhage: impact of CT on incidence. *Neurology* (NY), 1984; 34:653.
 68. Mohr JP et al: Stroke in south Alabama: incidence and diagnostic features — a population based study. *Stroke*, 1984; 15 :249.
 69. Brott T, Thalinger K: Hypertension as a risk factor for spontaneous intracerebral haemorrhage. *Stroke*, 1986; 17:1078.
 70. Drury I, Garraway WM : Primary intracerebral haemorrhage: impact of CT on incidence. *Neurology* (NY), 1984; 34:653.
 71. Gross CR, Kase CS et al: Stroke in south Alabama: incidence and diagnostic features — a population based study. *Stroke*, 1984; 15 :249.
 72. Tanaka H, Date C et al: Incidence of stroke in Shibata, Japan, 1976-1978. *Stroke*, 1981; 12:460.
 73. Ueda Y, Date C et al: Incidence of stroke in Shibata, Japan, 1976-1978. *Stroke*, 1981; 12:460.
 74. Furlan AJ, Whisnant JP: The decreasing incidence of primary intracerebral haemorrhage : a population study. *Ann Neurol*, 1979; 5:367.
 75. Elveback LR: The decreasing incidence of primary intracerebral haemorrhage : a population study. *Ann Neurol*, 1979; 5:367.
 76. Mohr JP, Caplan LR, Melski JW et al: The Harvard Cooperative Stroke Registry: a prospective registry. *Neurology* (NY), 1978; 28:754.
 77. Melski JW et al: The Harvard Cooperative Stroke Registry: a prospective registry. *Neurology* (NY), 1978; 28:754.
 78. Horsfield GI softening, with particular reference to heart weight. *J Pathol Bacteriol*, 1968; 96:311.
 79. Brewer DB, Fawcett FJ, Horsfield GI: A necropsy series of non-traumatic cerebral haemorrhages softening, with particular reference to heart weight. *J Pathol Bacteriol*, 1968; 96:311.
 80. Mutlu N, Berry RG, Alpers BG: Massive cerebral haemorrhage: clinical and pathological correlation. *Arch Neurol*, 1963; 8:74.
 81. McCormick WF, Rosenfield DB: massive brain haemorrhage: a review of 144 cases and an examination of their causes. *Stroke*, 1973; 4:946.
 82. McCormick, Schochet SS: *Atlas of cerebrovascular Disease*. P.328, WB Saunders, Philadelphia, 1976.
 83. Steering committee of the Physicians' Health study Research Group: Final report on the aspirin component of the Ongoing physicians' Health Study. *N Engl J Med*, 1989; 321:129.
 84. John Walton: *Brain's Diseases of the Nervous System* 12th edn 09:210-216.
 85. Lou H.C., Edvinsson L and MacKenzie, E.T (1987), The concept of coupling blood flow to brain function: revision required? *Ann Neurol*. 22, 289.
 86. Dolinskas CA, Belaniuk LT, Zimmerman RA, Kuhl DE: Computed Tomography of intracerebral hematomas. I. Transmission CT Observation on hematoma resolution. *AJR*, 1977; 129:681.
 87. Broderick J, Brott TG, Duldner JE et al: volume of intracerebral haemorrhage : a powerful and easy to use predictor of 30-day mortality. *Stroke*, 1993; 24:987.
 88. Siddharth N Shah. Clinical and Radiological profile of intra cerebral haemorrhage: *Journal of association of physicians of India* 2003; 51, 162.
 89. Siddharth N Shah. Clinical and CT evaluation of intracerebral haemorrhage: *journal of association of physicians of India* 2002, 50; 165:66.
 90. Abbott RD, Yin Y, Reed DM, Yanko K: Risk of stroke in Male cigarette smokers. *N Engl J Med*, 1986; 315:717.
 91. Donahue RP, Abbott RD, Reed DM, Yanko K: Alcohol and haemorrhagic stroke: The Honolulu Heart program. *JAMA*, 1986; 255:2311.
 92. Juvela S, Hillbom M, Palomaki H: Risk factors for spontaneous intracerebral haemorrhage. *Stroke*, 1995; 26:1558.
 93. Berry RG, Alpers BG: Massive cerebral haemorrhage: clinical and pathological correlation. *Arch Neurol*, 1963; 8:74.
 94. Belaniuk LT, Zimmerman RA, Kuhl DE: Computed Tomography of intracerebral hematomas. I. Transmission CT Observation on hematoma resolution. *AJR* 1, 1977; 29:681.
 95. Archana V, Reed DM, Yanko K: Risk of stroke in Male cigarette smokers. *N Engl J Med* 315:717, 1986.
 96. SK Mohapatra, Reed DM, : Alcohol and haemorrhagic stroke: The Honolulu Heart program. *JAMA* 255:2311, 1986.
 97. Akbari M, Palomaki H: Risk factors for spontaneous intracerebral haemorrhage. *Stroke* 26:1558, 1995.
 98. Amit Kumar, Onkar Nath Rai. *Study of risk factors among stroke patients in a tertiary hospital of Northern India*. *IJAM*. 2017, 2; 4:34-43.